

Age validation of canary rockfish (*Sebastes pinniger*) using two independent otolith techniques: lead-radium and bomb radiocarbon dating

Allen H. Andrews^{A,F}, Lisa A. Kerr^B, Gregor M. Cailliet^A, Thomas A. Brown^C, Craig C. Lundstrom^D and Richard D. Stanley^E

^AMoss Landing Marine Laboratories, California State University, 8272 Moss Landing Road, Moss Landing, CA 93940, USA.

^BUniversity of Maryland Center for Environmental Science, Chesapeake Biological Laboratory, PO Box 38, Solomons, MD 20688, USA.

^CLawrence Livermore National Laboratory, Center for Accelerator Mass Spectrometry, 7000 East Avenue, Livermore, CA 94551, USA.

^DUniversity of Illinois – Urbana Champaign, Department of Geology, 245 Natural History Building, 1301 West Green St, Urbana, IL 61801, USA.

^EFisheries and Oceans Canada, Pacific Biological Station, Nanaimo, British Columbia, V9T 6N7, Canada.

^FCorresponding author. Email: andrews@mlml.calstate.edu

Abstract. Canary rockfish (*Sebastes pinniger*) have long been an important part of recreational and commercial rockfish fishing from south-east Alaska to southern California, but localised stock abundances have declined considerably. Based on age estimates from otoliths and other structures, lifespan estimates vary from ~20 years to over 80 years. For the purpose of monitoring stocks, age composition is routinely estimated by counting growth zones in otoliths; however, age estimation procedures and lifespan estimates remain largely unvalidated. Typical age validation techniques have limited application for canary rockfish because they are deep-dwelling and may be long-lived. In this study, the unaged otolith of the pair from fish aged at the Department of Fisheries and Oceans Canada was used in one of two age validation techniques: (1) lead-radium dating and (2) bomb radiocarbon (¹⁴C) dating. Age estimate accuracy and the validity of age estimation procedures were validated based on the results from each technique. Lead-radium dating proved successful in determining that a minimum estimate of lifespan was 53 years and provided support for age estimation procedures up to ~50–60 years. These findings were further supported by $\Delta^{14}\text{C}$ data, which indicated that a minimum estimate of lifespan was 44 ± 3 years. Both techniques validate, to differing degrees, age estimation procedures and provide support for inferring that canary rockfish can live more than 80 years.

Additional keywords: accelerator mass spectrometry, age estimation, alpha-spectrometry, carbon-14, lead-210, radio-metry, radium-226, Scorpaenidae.

Introduction

Canary rockfish (*Sebastes pinniger*), a member of the family Scorpaenidae, is one of at least 65 species of rockfishes found in the north-eastern Pacific Ocean (Love *et al.* 2002). It ranges from south-eastern Alaska to northern Baja California, Mexico, with the centre of abundance from central California to British Columbia, Canada. The depth range extends from the surface to ~800 m, with most adults residing between 80 and 200 m (Love *et al.* 2002). The canary rockfish has long been an important part of the rockfish fishery with records of a commercial fishery off Washington and San Francisco, California dating to the 1880s (Love *et al.* 2002). It is typically caught in commercial trawl and hook-and-line fisheries from Monterey, California to British Columbia, Canada and is often taken with other members of

the genus in assemblages that are regionally defined (Yamanaka and Kronlund 1997; Bloeser 1999). The species is an important component of the recreational fishery throughout its range (except B.C., Canada), but declines in abundance led to significant harvest restrictions (i.e. *California Department of Fish and Game: Sportfishing Regulations and Pacific Fishery Management Council: 2003 Groundfish Management Regulations*). Canary rockfish was classified as overfished in 2000 based on a 1999 assessment (Methot and Piner 2002). A recent stock assessment of the canary rockfish fishery for the west coast of the USA indicated that the estimated depletion level of the spawning biomass in 2005 was between 6 and 11% (Methot and Stewart 2005); this is relative to estimates of unfished levels dating back to 1916 because of the long fishing history.

The targeted canary rockfish fishery off British Columbia, Canada is more recent; however, landings of rockfish were not recorded to species until the late 1960s and the Canadian stocks were likely exploited with other rockfishes as early as the 1950s (Stanley 1999). Landings have been consistent since 1967, but the status of Canadian stocks is poorly known. Management in British Columbia treats the canary rockfish fishery as two stocks: a central (Queen Charlotte Sound) and a southern (west coast of Vancouver Island) stock, with two-thirds of the landings traditionally coming from the southern stock. A documented decrease in mean age-at-capture based on the age composition of catches may indicate fishery exploitation has had a significant influence, but there are no convincing signs of over-exploitation. It is possible that stocks are close to maximum exploitation, but the actual status of the canary rockfish fishery for the waters of British Columbia is not well known (Stanley 1999).

Because management decisions are often made from age-based information, it is important to validate age and lifespan estimates as well as the age estimation procedures. Recent work using growth zone counting in otolith thin-sections provided estimates of lifespan for the canary rockfish from ~60 to more than 80 years (Chilton and Beamish 1982; Boehlert and Yoklavich 1986; Crone *et al.* 1999; Stanley 1999). Using the otolith burnt section technique (Fig. 1; MacLellan 1997), the Department of Fisheries and Oceans Canada (DFO) Ageing Laboratory in the Pacific Biological Station at Nanaimo, British Columbia has carried out routine canary rockfish age estimations for more than a decade and was responsible for the oldest estimated age of 84 years. Earlier studies estimated lifespan at little more than 20 years (Phillips 1964; Westheim and Harling 1975; Six and Horton 1977; Boehlert 1980), but this discrepancy can be explained by earlier studies using whole otoliths *v.* break-and-burn or sectioned otoliths (Boehlert and Yoklavich 1984). It is increasingly common to find that life-span estimates have increased for rockfishes with advances in otolith ageing and validation techniques (Cailliet *et al.* 2001).

Age validation techniques that can be applied to deep water fishes range widely in efficacy and age precision (Campana 2001). Some rely on establishing a temporal context to early growth by measuring changes in otolith zones or fish length (e.g. marginal increment analysis and length–frequency analysis); however, these techniques require an extrapolation of the age and growth information to older fish because of a loss of zone or length resolution. Other methods rely on marking and recapturing older fish (e.g. oxytetracycline injection and/or tagging), but deep-water, physoclistic fish usually have problems with survival upon return due to barotrauma and the techniques suffer because of low returns. Advances in the use of both naturally occurring and anthropogenic radioactive proxies for age have provided opportunities for independent age validation of deep-water fishes. The two primary techniques in use are lead-radium dating and the bomb radiocarbon (^{14}C) chronometer.

Lead-radium dating relies on the incorporation of naturally occurring radium-226 (^{226}Ra) from the environment into the otolith and its subsequent decay to lead-210 (^{210}Pb). By measuring the disequilibria of these two radioisotopes in otolith core material (first few years of life), an independent estimate of age can be determined based on the known ingrowth rate of ^{210}Pb from ^{226}Ra (Campana *et al.* 1990; Smith *et al.* 1991; Kimura




Fig. 1. Pictured is an example of an aged canary rockfish otolith using the burnt otolith section technique at the Department of Fisheries and Oceans Canada (DFO) Ageing Laboratory. Marked are the zones assumed to represent annual growth that are counted during routine age estimation. The estimated age of this section was 60 years.

and Kestelle 1995; Francis 2003). This technique works well as a tool for determining the validity of age interpretations that differ considerably, but its application is limited by relatively low resolution at ages approaching 100 years. The typical end result for this kind of study is that it provides strong support for a given age estimation technique and establishment of an independent estimate of lifespan that supports one age estimation technique over another. For example, lead-radium dating carried out on the yelloweye rockfish (*S. ruberrimus*) provided support for age exceeding 100 years (Andrews *et al.* 2002).

Use of the bomb ^{14}C chronometer requires a series of individual otoliths for which the birth years, based on standardised growth-zone age estimates, range from a time before significant atmospheric testing of thermonuclear devices (pre-1957) to the post-bomb period. This approach uses the discrete rise in ^{14}C from atmospheric testing as a time specific marker for

age validation. It is the agreement of the bomb ^{14}C record, from the species with age in question, with a reference bomb ^{14}C time-series that becomes a form of age validation. Hence, the usefulness of this approach for determining age or lifespan is dependent upon the difference between the collection year and time of first rise in ^{14}C for the reference time series. For example, a minimum lifespan of 30–38 years was determined for the red snapper (*Lutjanus campechanus*) based on these principles (Baker and Wilson 2001). In addition, age can also be determined by projecting measured ^{14}C levels back to a reference time series (Campana 1997; Andrews *et al.* 2005) or a series of samples with different collection years and ages can be selected to validate age estimation procedures for a range of age classes younger than the minimum lifespan (Piner and Wischniowski 2004).

Successful application of lead-radium dating and the bomb radiocarbon chronometer to rockfishes (family Scorpaenidae) has a well developed history. The first application of lead-radium dating to otoliths in general was an age validation study on the splitnose rockfish (*S. diploproa*) in which a 60-year lifespan was supported for the growth-zone age estimation criteria (Bennett *et al.* 1982). Since this pioneering application, the technique has been applied to several other rockfishes with varying degrees of age resolution (Campana *et al.* 1990; Kline 1996; Kestelle *et al.* 2000; Andrews *et al.* 2002, 2005; Stevens *et al.* 2004; Watters *et al.* 2006). For the bomb radiocarbon technique, its application to fish otoliths began with fishes in the southern hemisphere (Kalish 1993, 2001), followed by a series of studies in the northern hemisphere (e.g. Campana 1997; Kalish *et al.* 2001).

The first application of the bomb ^{14}C technique to rockfishes was a follow-up study to age validation work on yelloweye rockfish (Andrews *et al.* 2002). Because yelloweye rockfish were relatively easy to age from growth zone counts (coefficient of variation [CV] = 4%) and lead-radium dating provided support for the age estimation procedures, a series of yelloweye rockfish otoliths were used to establish a bomb radiocarbon chronometer for the north-eastern Pacific Ocean (Kerr *et al.* 2004). The chronometer established by Kerr *et al.* (2004) was further corroborated by an independent study of juvenile Pacific halibut (*Hippoglossus stenolepis*) from a similar region (Piner and Wischniowski 2004). These records subsequently provided an independent chronometer for age validation of the quillback (*S. maliger*; Kerr *et al.* 2005) and bocaccio (*S. paucispinis*; Andrews *et al.* 2005) rockfishes. A recent application of this technique to canary rockfish collected off Oregon, and aged by the Oregon Department of Fish and Wildlife, provided unexpected results that were attributed to some combination of age estimate bias and regional oceanography (Piner *et al.* 2005).

The purpose of this study was to assess the validity of age estimates and age estimation procedures used in routine age assignments of canary rockfish by the DFO, along with any agency that uses the same age estimation criteria (MacLellan 1997). Lead-radium dating and bomb ^{14}C analyses will be used to provide a basis for support or dispute of canary rockfish age estimates determined using the break and burn method. These results will also be compared with recent findings from a canary rockfish age validation study for collections made off Oregon (Piner *et al.* 2005). Additionally, variation in regional pre-bomb ^{14}C values for the north-eastern Pacific Ocean was examined

along with the potential influence of this variability on age validation.

Materials and methods

The high estimated lifespan and the availability of a well developed otolith archive at DFO made the canary rockfish an excellent candidate for a combined application of the two independent age validation techniques. Otoliths were selected from these archives based on the criteria for each technique. The lead-radium dating required groups of otoliths and the bomb radiocarbon technique required only a series of carefully chosen otoliths with estimated birth dates spanning the pre-bomb to post-bomb period. The application of each technique was treated as an independent project and is separated into sections for clarity.

Lead-radium dating

Otoliths were selected for analysis from more recent years between 1990 and 1998 based on the number of aged fish available. Age groups were chosen to cover a wide range of estimated age and the groups were as close to a single age as possible. The size of the core material extracted was determined based on 33 whole juvenile otoliths aged at 3 years. Average size of the otoliths was 9.0 mm L \times 4.8 mm W \times 1.2 mm D and the average weight was 0.073 ± 0.012 (s.d.) g. This target core size could be easily extracted and consisted of just a few years of growth to minimise the possible error associated with variable ^{226}Ra uptake (Francis 2003). Otoliths from adult fish were ground down to the target core size using a lapping wheel and 80–120 grit silicon-carbide paper. The radiometric analysis of the pooled otolith core samples for ^{210}Pb and ^{226}Ra followed procedures previously described in Andrews *et al.* (1999a, 1999b, 2005).

Measured $^{210}\text{Pb} : ^{226}\text{Ra}$ activity ratios for each age group, along with total sample age (growth-zone age plus time since capture), were plotted with the $^{210}\text{Pb} : ^{226}\text{Ra}$ ingrowth curve to provide a reference to the ratio that would be expected for a given age. The uncertainty for the growth-zone derived age of each group was represented by the age estimate CV (5.8%), as was determined from the variability of age estimates among three readers at DFO. This uncertainty was assigned to the mean age of each group as some indication of what might be the limit for the degree of precision in the age estimation criteria, and should not be confused with the degree of accuracy. Agreement between radiometric age and growth zone derived age was determined by comparing the limits of 95% confidence with the ingrowth curve. To provide a lower limit on the minimum lifespan for the fish aged in this study, the 95% CI from the $^{210}\text{Pb} : ^{226}\text{Ra}$ ratios provided a validated basis for such a determination.

Bomb radiocarbon

Otoliths from aged canary rockfish were selected from the DFO archive based on estimated birth date to cover the pre-bomb to post-bomb period. Most of the fish specimens that provided the otoliths were captured off the west coast of Vancouver Island. Otoliths used in the radiocarbon part of this study were from three 'Major Areas' used for landings record keeping off British Columbia by DFO (3C–D and 5A–B). The most important samples for recording the rise in ^{14}C (1950–1967) were all from

Table 1. List of attributes for each canary rockfish sample (age group) collected off British Columbia, Canada that was analysed using lead-radium dating

Sample number	Average age and range (year)	Fork length and range (cm)	Average collection date and range	Core number Male/Female
CR1	3 (3)	19.1 (19–21)	5/91	16 ^A
CR2	10 (10)	38.6 (33–46)	6/21/91 (6/14/91–7/15/91)	8/10
CR3	18.8 (18–20)	53.2 (44–58)	3/21/91 (2/13/91–6/14/91)	11/9
CR4	52.2 (51–54)	53.7 (51–56)	12/12/94 (7/14/93–10/14/95)	20/0
CR5	52.0 (50–54)	52.5 (51–55)	9/8/90 (4/14/90–11/13/90)	6/0
CR6	11 (11)	45.0 (40–48)	3/5/91 (2/13/91–3/15/91)	3/3
CR7	41.0 (40–42)	54.0 (49–57)	3/19/98 (2/12/98–4/14/98)	8/0
CR8	45.3 (45–46)	54.1 (52–59)	3/28/98 (3/15/98–4/14/98)	5/2
CR9	55.4 (55–56)	53.2 (50–55)	3/15/98 (2/12/98–4/14/98)	5/0
CR10	3 (3)	19.9 (19–21)	5/91	16 ^A
CR11	51.8 (50–53)	55.2 (52–57)	3/15/98 (2/12/98–4/14/98)	6/0
CR12	53.6 (53–54)	54.4 (52–57)	5/3/98 (2/12/98–11/13/98)	4/1

^AImmature fish.

Major Area 3D. The other samples were from Major Areas nearby: 1945 (5A), 1969 and 1973 (5B), and 1980 (3C).

The core of each otolith, which constituted less than the first year of growth, was analysed for $\Delta^{14}\text{C}$ because it is known that at least the first year's growth was formed while the fish inhabited the ocean mixed layer. The bomb radiocarbon technique is most reliable for fishes that inhabit the mixed layer of the ocean, at least during a portion of their life history, because of uncertainty regarding mixing rate at depth and limited data on the $\Delta^{14}\text{C}$ signal in deeper waters (Kalish 2001).

The size of the core material extracted was determined from nine whole juvenile otoliths with an age of one year. Average dimensions were 6.9 mm L \times 3.8 mm W \times 0.9 mm D with a weight of \sim 0.02 g. Sampling within these dimensions was targeted to avoid the inclusion of material formed later in life. Core extraction was carried out using a milling machine with a 3.2 mm (1/8") diameter end mill, pressed into the otolith to a depth of 0.8 mm. Visual alignment of the milling machine bit with the centre of the distal side of the otolith allowed for correction due to individual variability; the first year's growth was easily recognisable because more recent accretion on the distal surface was negligible. Coring produced small powdered samples that were weighed to the nearest 0.1 mg. For $\Delta^{14}\text{C}$ analysis, otolith calcium carbonate (CaCO_3) was converted to graphite and measured for ^{14}C content on an accelerator mass spectrometer (AMS) at the Center for Accelerator Mass Spectrometry, Lawrence Livermore National Laboratory, Livermore, California, USA. The ^{14}C measurements were reported as $\Delta^{14}\text{C}$ values and a measured $\delta^{13}\text{C}$ value of -3.1 , similar to marine carbonates, was used in calculating the $\Delta^{14}\text{C}$ values (Stuiver and Polach 1977).

The $\Delta^{14}\text{C}$ values measured in the canary rockfish otolith cores were plotted with respect to the estimated birth year, which was estimated based on the difference between estimated age and the collection year. For each birth year a range of uncertainty between readers was calculated based on the age estimation CV (5.8%). This uncertainty was assigned to the birth years as some indication of what might be the limit for the degree of precision in the age estimation criteria, and should not be confused with the degree of accuracy. Agreement of the estimated birth

year range with the reference time-series, coupled with use of the exponential fit method, as suggested in Kerr *et al.* (2004), was used as a basis for providing a minimum lifespan and age estimate accuracy. Specifically, the first sample below the two s.d. criterion of pre-bomb $\Delta^{14}\text{C}$ levels was used to determine a minimum lifespan. The reference-time series used for comparisons were from yelloweye rockfish (*S. ruberrimus*: Kerr *et al.* 2004) and Pacific halibut (*Hippoglossus stenolepis*: Piner and Wischniowski 2004) because of the regional similarities to the canary rockfish collection locations and specifically yelloweye rockfish for the similarities in life history.

Results

Lead-radium dating

Twelve samples, representing groups of otoliths, were analysed from fish that ranged in the growth-zone derived age from 3 to over 50 years (Table 1). Typically, numerous otoliths are required to attain enough material for measurable ^{210}Pb and ^{226}Ra from the age groups, resulting in a wide age range for ages with low sample availability. For the first set of four samples (CR1–CR4) the target number was \sim 20 otolith cores because levels of ^{210}Pb and ^{226}Ra were unknown. Once preliminary findings indicated less material was needed, fewer otolith cores were pooled to narrow the estimated age range for the rest of the samples. Fish length ranged from \sim 20 cm FL for the juvenile groups (CR1 and CR10) to more than 50 cm FL for the older age groups. Collection years varied considerably (1990–1998) because otolith age groups were selected based on the number of otoliths available and narrowness of age range for the groups. For most groups the collection dates were all within a few months of each other. Sexes were not separated because validation of the age estimation method was the goal. Total numbers of otoliths per age group ranged from five to twenty and most were male.

Radiometric analysis of all age groups ($n = 12$) resulted in the successful determination of ^{210}Pb activities, and limited success for ^{226}Ra activities (Table 2). Activities of ^{210}Pb were variable as expected and typically increased with estimated age. No recovery of ^{226}Ra in the first set of four samples was due to a mistake

Table 2. Radiometric results for canary rockfish sample series

Listed are the estimated age range from growth-zone counts and measured ^{210}Pb and ^{226}Ra activities for all samples, except where ^{226}Ra was not recovered (N.R.). Two standard errors were expressed as an error percentage for the measured activities. Calculated activity ratios and the corresponding margin of error (based on error propagation and the delta method) are given to provide a 95% confidence interval

Sample number	Age group (year)	$A^{210}\text{Pb}$ (dpm g^{-1}) \pm % error	$A^{226}\text{Ra}$ (dpm g^{-1}) \pm % error	$^{210}\text{Pb} : ^{226}\text{Ra}$ activity ratio
CR1	3	0.0381 ± 6.9	N.R.	N.A. ^A
CR2	10	0.0377 ± 6.2	N.R.	N.A. ^A
CR3	18–20	0.0760 ± 4.9	N.R.	N.A. ^A
CR4	51–54	0.0727 ± 5.3	N.R.	N.A. ^A
CR5	50–54	0.0797 ± 7.2	0.0899 ± 6.4	0.866 (0.085)
CR6	11	0.0478 ± 8.5	0.102 ± 4.3	0.470 (0.045)
CR7	40–42	0.0806 ± 6.8	0.106 ± 13.9	0.761 (0.118)
CR8	45–46	0.0942 ± 6.7	0.109 ± 8.0	0.863 (0.090)
CR9	55–56	0.0632 ± 7.8	0.0924 ± 8.6	0.684 (0.080)
CR10	3	0.0396 ± 9.5	0.122 ± 6.1	0.326 (0.037)
CR11	50–53	0.0726 ± 7.7	0.0803 ± 1.3	0.904 (0.071)
CR12	53–54	0.0707 ± 7.4	0.0888 ± 3.8	0.795 (0.066)

^ANot applicable (N.A.) because the activity of ^{226}Ra was highly variable and precluded use of an average.




Fig. 2. Plot of the measured $^{210}\text{Pb} : ^{226}\text{Ra}$ ratios with respect to total sample age (growth-zone age plus the time since capture) for canary rockfish samples, plotted with the $^{210}\text{Pb} : ^{226}\text{Ra}$ ingrowth curve. Horizontal error bars (some within the symbol size) represent age estimate uncertainty (coefficient of variation [CV] = 5.8%) relative to the mean growth-zone age. The vertical error bars represent the analytical uncertainty associated with measuring ^{210}Pb and ^{226}Ra (two s.e.). Dashed lines below signify the affect of age estimate bias for over estimation of age ($b = 25\%$ and $b = 50\%$).

in the mixing of acid that led to changes in the elution characteristics and a loss of radium. Radium was recovered for all other samples ($n = 8$) and resulted in a variable range of activity (range of $0.0803\text{--}0.122$ dpm g^{-1} (disintegrations per minute per gram)). This degree of variability precluded the determination of a radiometric age for the samples where radium was lost because levels were not consistent enough to use an average ($0.0988 \pm 11\%$ 2SE). Although the activity levels for both ^{210}Pb and ^{226}Ra were quite variable, the $^{210}\text{Pb} : ^{226}\text{Ra}$ activity ratio increased as expected with growth-zone derived age. The

importance of sample-specific ^{226}Ra activity for this species was exemplified by the 3-year age group (sample CR10), which had the highest measured ^{226}Ra activity in this study and, as expected because of the young age, the lowest $^{210}\text{Pb} : ^{226}\text{Ra}$ ratio.

The measured $^{210}\text{Pb} : ^{226}\text{Ra}$ ratios were largely in agreement with the ingrowth curve (Fig. 2). All but two samples (CR9 and CR11) were in agreement with growth-zone derived age based on the 95% CI from the measured ratios. To give an indication of the affect of growth-zone ageing bias, the plot included two ingrowth curves that incorporate an age overestimation bias of

25% and 50%. The agreement between growth-zone derived age and $^{210}\text{Pb} : ^{226}\text{Ra}$ ratios provide support for the age estimates and age estimation procedures.

Radiometric ages determined for the eight samples from which radium was recovered ranged from 3 to 71 years (Table 3). These ages were corrected for time since collection based on the difference from the date of processing and average collection date for the age group. Greater radiometric age uncertainty for the oldest samples was because of the asymptotic nature of the ingrowth curve; small changes in the measured ratio become large changes in radiometric age as the ratio approaches 1.0. By treating lead-radium dating as an independent estimate of age, there was 95% confidence that the oldest canary rockfish used in this study were at least 53 years of age (CR11).

Bomb radiocarbon

Selecting the fish for ^{14}C analysis via AMS resulted in twelve canary rockfish otoliths with estimated birth dates ranging from

Table 3. Summary of growth-zone ages and radiometric ages for canary rockfish age groups

Average growth-zone age (CV of 5.8% relative to the mean) are given with radiometric age calculated from the measured $^{210}\text{Pb} : ^{226}\text{Ra}$ activity ratios and corrected for time since capture. Radiometric age range was based on the analytical uncertainty and error propagation (two s.e.)

Sample number	Age group (year)	Average growth-zone age (year)	Radiometric age range (year)
CR5	50–54	52 (2.9)	59 (41–103)
CR6	11	11 (0.6)	10 (8–13)
CR7	40–42	41 (2.3)	43 (30–65)
CR8	45–46	45 (2.5)	61 (44–95)
CR9	55–56	55 (3.1)	34 (27–43)
CR10	3	3 (0.1)	3 (1–4)
CR11	50–53	52 (2.9)	71 (53–114)
CR12	53–54	54 (3.0)	47 (38–60)

1945 to 1980 (Table 4). Collection years ranged from 1991 to 1998, with age estimates of 11–49 years. The choice of available otoliths was designed to have samples with a range of birth years that encompassed the period of ^{14}C increase (pre-bomb to post-bomb). Extracted mass of core material ranged from 3.6 to 7.3 mg – well below the weight of a one-year juvenile otolith. The measurement of $\Delta^{14}\text{C}$ from this series of otolith cores ranged from -113.1 ± 4.6 to $81.9 \pm 3.8\%$ for the birth years 1945 and 1967, respectively.

Plotting the $\Delta^{14}\text{C}$ data with the two reference time series (yelloweye rockfish and Pacific halibut) revealed a strong conformance for the canary rockfish data (Fig. 3). Pre-bomb $\Delta^{14}\text{C}$ levels for the series of samples with estimated birth years 1945–1956 had an average of $-105 \pm 8\%$, similar to the pre-bomb time series for the yelloweye rockfish ($-102 \pm 9\%$) and Pacific halibut ($-107 \pm 7\%$). The time of first rise in $\Delta^{14}\text{C}$ for the canary rockfish record appears to be between 1957 and 1961, or as late as 1963, as was subjectively determined by visual examination of the plot. Application of the exponential fit method provided an objective means of determining the year of initial rise from pre-bomb levels. The fitted curve ($y = -115.06 + 1.414E-154 \cdot \exp[0.183 \cdot X]$) and time of intersection with the two s.d. criterion ($\sim \Delta^{14}\text{C} = -89$) indicated the first rise in ^{14}C was 1958. This finding was in agreement with both reference time series and more typical of the yelloweye rockfish distribution. The rapid increase in ^{14}C from 1961 ± 2 years to a maximum at 1967 ± 2 years is also in agreement with both reference time series. The peak and post-bomb $\Delta^{14}\text{C}$ levels of the canary rockfish more closely resemble the yelloweye rockfish distribution, although the peak $\Delta^{14}\text{C}$ level was greater.

A minimum age or lifespan was determined based on the difference between collection year and the estimated birth year for the last sample with pre-bomb $\Delta^{14}\text{C}$ levels. Because the year of first rise in ^{14}C was calculated as 1958 at a $\Delta^{14}\text{C}$ level of -89% , the last sample to contain levels below this value was the sample with an estimated birth year of 1954 ± 3 years

Table 4. Summary of fish age and otolith data for canary rockfish collected off British Columbia, Canada

Resolved age was the final age estimate assigned by the Department of Fisheries and Oceans Canada (DFO). Birth year was the collection year minus the resolved age. Age error was the uncertainty associated with the age estimate (CV = 5.8%). Radiocarbon values from the otolith cores of canary rockfish were expressed as $\Delta^{14}\text{C}$ with the accelerator mass spectrometer (AMS) analytical uncertainty (s.d.)

Resolved age (year)	Collection year	Birth year ($\pm 5.8\%$ CV)	Core sample weight (mg)	$\Delta^{14}\text{C}$ (‰) \pm s.d.
49	1994	1945 ± 3 years	5.8	-113.1 ± 4.6
48	1998	1950 ± 3 years	3.6	-97.3 ± 3.1
44	1998	1954 ± 3 years	5.2	-104.5 ± 3.2
41	1998	1957 ± 3 years	7.3	-88.6 ± 3.2
38	1998	1960 ± 2 years	6.3	-96.4 ± 3.3
37	1998	1961 ± 2 years	4.4	-84.3 ± 3.1
35	1998	1963 ± 2 years	8.6	-61.8 ± 3.2
33	1998	1965 ± 2 years	4.4	50.2 ± 4.6
31	1998	1967 ± 2 years	5.6	81.9 ± 3.8
22	1991	1969 ± 1 years	5.1	62.0 ± 3.8
18	1991	1973 ± 1 years	5.4	34.6 ± 3.6
11	1991	1980 ± 1 years	6.0	18.0 ± 3.8




Fig. 3. Plot of measured $\Delta^{14}\text{C}$ values for the canary rockfish otoliths with respect to estimated birth year, derived from growth zone counts determined by the Department of Fisheries and Oceans Canada (DFO). The horizontal bars represent the range of possible birth dates from the age estimate uncertainty (coefficient of variation [CV] = 5.8%). Note the strong conformity of the canary rockfish data with the yelloweye rockfish and Pacific halibut reference time series.

(-104.5 ± 3.2). Based on the difference from collection year (1998) the age of this fish was at least 39 ± 2 years and the findings support a break-and-burn based age estimate of 44 ± 3 years for this sample. In addition, assuming the rise of bomb-produced $\Delta^{14}\text{C}$ levels up to peak values can also be used as a potential measure of bias, it was further concluded that there was no evidence for ageing bias that was greater than the age estimate precision for the fish aged 31 ± 2 to 38 ± 2 years.

Discussion

Lead-radium dating

The lead-radium dating sample series illustrated a close conformance to the expected lead-radium ingrowth curve with minor exceptions. These data provided general support for ages up to ~ 50 – 60 years based on the trend. Because some of the old age groups were approaching the asymptote of the ingrowth curve, small changes in the $^{210}\text{Pb} : ^{226}\text{Ra}$ ratio resulted in a higher degree of uncertainty. While the upper limit was not well defined, there was 95% confidence that the oldest canary rockfish used in this study were at least 53 years of age. These findings provide support for (1) canary rockfish age estimation procedures used by DFO, and (2) for lifespan estimates exceeding 80 years based on a continued adherence to the age estimation protocol.

The measured $^{210}\text{Pb} : ^{226}\text{Ra}$ activity ratios conformed well to the ingrowth curve for total sample age. Note that the total age of the 3-year age group was in close agreement with the expected $^{210}\text{Pb} : ^{226}\text{Ra}$ activity ratio for a total age of 14 years. This is an indication that the extracted otolith continues to function as a closed system after removal from the fish and that the initial uptake of exogenous ^{210}Pb was negligible. Findings of this kind, coupled with studies directed at testing for intermediate isotope loss or exogenous incorporation of ^{210}Pb (i.e. Baker *et al.* 2001; Kastle and Forsberg 2002), continue to refute concerns about

violations to the closed system assumption (e.g. Gauldie and Cremer 2000).

Bomb radiocarbon

The conformity of the canary rockfish ^{14}C record in the present study with the yelloweye rockfish and Pacific halibut $\Delta^{14}\text{C}$ reference time-series was compelling information that supported the age estimates and current age estimation procedures. Considering the similarity of biology, ecology, and distribution of the two rockfishes, the use of the yelloweye rockfish $\Delta^{14}\text{C}$ time-series as a means of temporal calibration for the canary rockfish record was well supported. Unlike the quillback rockfish, which has a more nearshore distribution as larvae, the canary rockfish $\Delta^{14}\text{C}$ record for the period beyond the post-bomb peak was also similar to the yelloweye rockfish time-series. The timing of the initial rise in $\Delta^{14}\text{C}$ for canary rockfish otoliths (1957 ± 2 years) was consistent with the yelloweye rockfish record (1958 ± 2 years). In addition, it was further concluded that there was no evidence for ageing bias that was greater than the age estimate precision for fish aged 31 ± 2 to 38 ± 2 years because no phase shift was documented in the samples aged as having birth years between the first rise to peak of $\Delta^{14}\text{C}$ levels. It is reasonably certain that the samples with pre-bomb $\Delta^{14}\text{C}$ levels in core material from otoliths collected in 1998 were at least 39 ± 2 years and that a minimum estimate of lifespan was 44 ± 3 years was supported by the measured $\Delta^{14}\text{C}$ in the first pre-bomb sample.

Regional pre-bomb $\Delta^{14}\text{C}$ levels in the mixed layer of the ocean are assumed to be relatively constant; however, variation in these levels may introduce uncertainty into age validation studies because baseline levels provide the reference for the time of first rise. Pre-bomb levels of ^{14}C and the degree of variation were similar between the canary rockfish and the reference time series (yelloweye rockfish and Pacific halibut) and considered together

(samples with birth dates before 1957) provide a mean pre-bomb $\Delta^{14}\text{C}$ value of $-105 \pm 8\%$ ($n = 23$). Adding the results from the quillback rockfish study (Kerr *et al.* 2005) provided a grand mean $\Delta^{14}\text{C}$ of $-104 \pm 10\%$ ($n = 26$) for the Gulf of Alaska region. This value included samples ranging from the west coast of Vancouver Island, British Columbia, Canada to the northern Gulf of Alaska. To consider this value a characteristic of the regional $\Delta^{14}\text{C}$ average during the pre-bomb period (est. 1940–1956 from the range of estimated birth years), the biggest assumption is some degree of regional specificity for the ontogeny of each fish sampled. For the Pacific halibut samples, five samples were from young of the fish in the year that experienced a pelagic phase of up to six to seven months and six samples were 3–5-year fish (cored to the first year) that would have experienced the pelagic phase and possibly a migration period (Piner and Wischniowski 2004). Rockfishes are known to have a pelagic larval stage as well, but the potential complication for using these data as regional indicators of ^{14}C is potential movement over great distances as an adult.

Ontogenetic movement of rockfishes varies considerably among species; many are relatively residential and some can travel great distances (see Love *et al.* 2002 for a synopsis). The species considered here are not known to move great distances. Tagged yelloweye rockfish off Depoe Bay were residential while at liberty for up to 739 days; however, of the 23 tagged canary rockfish returns off Oregon some were residential and 14 exhibited movement of more than 100 km north or south to a maximum of 236 km from the release location (DeMott 1983; Bill Barss, Oregon State University [*emeritus*], per. comm.). This degree of movement, given that it may apply to the other species considered here, could broaden the apparent range for the regional mean pre-bomb $\Delta^{14}\text{C}$ levels exhibited by these species.

The level of natural pre-bomb $\Delta^{14}\text{C}$ ranged from -76.9 to -120.0% , for the four species of Gulf of Alaska fishes. This level of variation could affect the temporal distribution of the bomb signal. An example of how low $\Delta^{14}\text{C}$ values can be in the nearshore waters of a region was from pre-bomb levels that ranged well below -120% off Adak Island, Alaska as evidenced by the study of an alga (*Clathromorphum nereostratum*) that resided at a depth of 10 m (Frantz *et al.* 2005). A deep-water ^{14}C record measured from red tree coral (*Primnoa pacifica*, syn. *P. resedaeformis*) from 263 m in the Dixon Entrance, south-eastern Alaska indicated that levels can be quite depleted (-129.8 to -136.4% ; A. Andrews, unpub. data). Thus, deep-water sources advected to nearshore waters can affect the shape and apparent timing of the first rise in $\Delta^{14}\text{C}$, resulting in a vertical attenuation of the bomb signal. This factor provides a reasonable explanation for what appears to be a phase lag of a few years for some samples in the yelloweye and canary rockfish records, but can also be explained by small variations in age interpretation, as is indicated by the error bars.

A recent study, which also used bomb $\Delta^{14}\text{C}$ as a tool for age validation, of canary rockfish collected off Oregon provided results that appear to differ from the current study in the timing and magnitude of ^{14}C rise (Piner *et al.* 2005). Pre-bomb levels off Oregon were similar to the levels reported in this study ($-105 \pm 8\%$), based on four samples with estimated birth years ranging from 1949 to 1956 ($\sim -103 \pm 4\%$); however,

post-bomb levels appear to have increased more gradually and many samples appear to have a significant phase lag of up to 7 years (average 2–3 years). In addition, the post-bomb peak values appear to be much lower than recorded for the canary rockfish taken off Canada (30.0 ± 3.5 cf. $78.5 \pm 4.3\%$). As stated above with regard to natural variability in pre-bomb $\Delta^{14}\text{C}$ levels, it is possible that some of the phase lag in the Oregon canary rockfish study was due to this effect, but it is unlikely that it could be responsible for the low values with rather large apparent phase lags. For example, an apparent lag of 7 years relative to the time of first rise in $\Delta^{14}\text{C}$ would require depressions of peak $\Delta^{14}\text{C}$ levels in the order of 100%.

Further examination of the data from the Oregon canary rockfish study revealed that age interpretation alone can explain most of the unexpected distribution, as was discussed by Piner *et al.* (2005), who listed two or three age estimates for each sample. They chose to use the first readings in their analyses because they were made following methodologies used in stock assessments (Crone *et al.* 1999; Methot and Piner 2002). However, the second and/or third readings tended to be older (with differences up to 4 years), and when the oldest age determinations were used many of the critical samples in the unexpected distribution were shifted further towards what would be expected in comparison to the yelloweye rockfish and Pacific halibut records (e.g. highest $\Delta^{14}\text{C}$ value at 1967 cf. 1970; see Fig. 4). Hence, it is likely that the timing of the rise in $\Delta^{14}\text{C}$ was closer to 1959–1962 than was depicted in Piner *et al.* (2005). This finding implies that the later otolith readings were more accurate and an inter-calibration between DFO and ageing laboratories in the USA is recommended. It can be surmised that the study validated a level of age estimate imprecision to an average of ~ 2.8 years based on the observed phase shift (Piner *et al.* 2005). When considering this level of age estimate imprecision, it is also possible that the unexpectedly low $\Delta^{14}\text{C}$ peak value ($30.0 \pm 3.5\%$) can be attributed to a lack of fish young enough to capture the regional peak $\Delta^{14}\text{C}$ level.

Given that pre-bomb levels were not more depressed for the Oregon canary rockfish record, it is unlikely that oceanography off Oregon was drastically different in terms of $\Delta^{14}\text{C}$ circulation from regions of upwelling further north in the Gulf of Alaska. In support of this hypothesis, the range of pre-bomb $\Delta^{14}\text{C}$ values available from all fish records for the north-eastern Pacific Ocean was -76.9 to -120.0% with an overall average of $-103 \pm 9\%$ (Fig. 5). In addition, a recent study of cowcod (*Sebastes levis*) otoliths off Southern California provided a measure of pre-bomb levels much further south with a mean of $-97 \pm 3\%$ ($n = 4$; A. Andrews, unpub. data). Less depleted $\Delta^{14}\text{C}$ values for more southerly coastal waters of the north-eastern Pacific Ocean have been attributed to less influence from upwelling (Ingram and Southon 1996). The unexpectedly low levels for post-bomb measurements in the study of canary rockfish off Oregon are difficult to explain in these terms and are more suggestive of age estimation problems.

Conclusions

The findings of this study provided robust support for canary rockfish age estimates. Lead-radium dating proved successful by showing conformance to the expected lead-radium ingrowth




Fig. 4. Plot of data taken from Piner *et al.* (2005) for canary rockfish collected off Oregon with data from this study and the yelloweye rockfish reference time-series. The birth dates for the Oregon canary rockfish, in this case, have been plotted with alternate birth years calculated from other age interpretations listed in the study (Piner *et al.* 2005). This figure is provided to illustrate that the findings for Oregon canary rockfish may have been better than initially interpreted and that the second and third otolith readings in the study were probably more accurate.




Fig. 5. Average pre-bomb $\Delta^{14}\text{C}$ from fish otolith records throughout the north-eastern Pacific Ocean. Species and associated values are placed central to the region that samples were taken. All samples were from coast and shelf waters along the Gulf of Alaska to the California coast. The value for Pacific halibut is placed as such because the samples ranged from off Kodiak Island, Alaska to Vancouver Island, British Columbia, Canada.

curve and provided support for ages up to ~50–60 years. At the very least, there was 95% confidence that lead-radium dating determined a minimum lifespan of 53 years, discounting early lifespan estimates on the order of 20 years. These findings were further supported by $\Delta^{14}\text{C}$ data that indicated canary rockfish have a minimum lifespan of at least 44 ± 3 years. In addition, the conformity of $\Delta^{14}\text{C}$ measurements from otolith cores with estimated birth years for the post-bomb era provided further support for these age estimates down to ~30 years. To differing degrees, both techniques provided support for the age estimation procedures used in routine age assignments for canary rockfish by the DFO Ageing Laboratory and an age validated basis for inferring that canary rockfish can live more than 80 years.

The investigation of pre-bomb $\Delta^{14}\text{C}$ levels in the mixed layer of the Gulf of Alaska from regional fish otolith records resulted in a mean $\Delta^{14}\text{C}$ of $-104 \pm 10\%$ ($n = 26$) with a potential range of -76.9% to -120.0% . This relatively high degree of regional variability in pre-bomb levels would have had a natural effect on the apparent timing of the first rise in $\Delta^{14}\text{C}$ and may provide a reasonable explanation for what appears to be a phase lag of a few years for some samples. When considering all fish studies ranging from Southern California to the Gulf of Alaska, including cowcod and the five oldest Oregon canary rockfish samples (Piner *et al.* 2005), a grand mean $\Delta^{14}\text{C}$ of $103 \pm 9\%$ ($n = 35$) was calculated. Based on this observation the levels for pre-bomb $\Delta^{14}\text{C}$ in the mixed layer were fairly constant in a much broader regional context.

Acknowledgements

We thank Shayne MacLellan and Darlene Gillespie of the Department of Fisheries & Oceans Canada, Pacific Biological Station, Nanaimo, B.C., Canada for providing aged canary rockfish otoliths for this study in addition to assistance with manuscript preparations. Thanks to Melissa Stevens for assisting with processing the otolith samples and age groups analysed in this study, Paula Zermeño at the Center for Accelerator Mass Spectrometry for assistance with the bomb radiocarbon analyses, and Kenneth Coale for assistance with data interpretation. This work was supported by the National Sea Grant College Program of the USA Department of Commerce's National Oceanic and Atmospheric Administration under NOAA Grant number NA06RG0142, project numbers R/F-182 and R/F-190, through the California Sea Grant College Program; and in part by the California State Resources Agency. This research was also supported by the Center for Accelerator Mass Spectrometry under the University Collaborative Research Program at Lawrence Livermore National Laboratory and was carried out, in part, under the auspices of the USA Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48. The views expressed herein do not necessarily reflect the views of any of those organisations. The USA government is authorised to reproduce and distribute for governmental purposes.

References

- Andrews, A. H., Cailliet, G. M., and Coale, K. H. (1999a). Age and growth of the Pacific grenadier (*Coryphaenoides acrolepis*) with age estimate validation using an improved radiometric ageing technique. *Canadian Journal of Fisheries and Aquatic Sciences* **56**, 1339–1350. doi:10.1139/CJFAS-56-8-1339
- Andrews, A. H., Coale, K. H., Nowicki, J. L., Lundstrom, C., Palacz, Z., Burton, E. J., and Cailliet, G. M. (1999b). Application of an ion-exchange separation technique and thermal ionization mass spectrometry to ^{226}Ra determination in otoliths for radiometric age determination of long-lived fishes. *Canadian Journal of Fisheries and Aquatic Sciences* **56**, 1329–1338. doi:10.1139/CJFAS-56-8-1329
- Andrews, A. H., Cailliet, G. M., Coale, K. H., Munk, K. M., Mahoney, M. M., and O'Connell, V. M. (2002). Radiometric age validation of the yelloweye rockfish (*Sebastes ruberrimus*) from southeastern Alaska. *Marine and Freshwater Research* **53**, 139–146. doi:10.1071/MF01126
- Andrews, A. H., Burton, E. J., Kerr, L. A., Cailliet, G. M., Coale, K. H., Lundstrom, C. C., and Brown, T. A. (2005). Bomb radiocarbon and lead-radium disequilibria in otoliths of bocaccio rockfish (*Sebastes paucispinis*): a determination of age and longevity for a difficult-to-age fish. Proceedings of the 3rd International Symposium on Otolith Research and Application. *Marine and Freshwater Research* **56**, 517–528. doi:10.1071/MF04224
- Baker, M. S., and Wilson, C. A. (2001). Use of bomb radiocarbon to validate otolith section ages of red snapper *Lutjanus campechanus* from the northern Gulf of Mexico. *Limnology and Oceanography* **46**, 1819–1824.
- Baker, M. S., Jr, Wilson, C. A., and VanGent, D. L. (2001). Testing assumptions of otolith radiometric aging with two long-lived fishes from the northern Gulf of Mexico. *Canadian Journal of Fisheries and Aquatic Sciences* **58**, 1244–1252. doi:10.1139/CJFAS-58-6-1244
- Bennett, J. T., Boehlert, G. W., and Turekian, K. K. (1982). Confirmation of longevity in *Sebastes diploproa* (Pisces: Scorpaenidae) from $^{210}\text{Pb}/^{226}\text{Ra}$ measurements in otoliths. *Marine Biology* **71**, 209–215. doi:10.1007/BF00394632
- Bloeser, J. A. (1999). 'Diminishing Returns: the Status of West Coast Rockfish.' (Pacific Marine Conservation Council: Astoria, OR.)
- Boehlert, G. W. (1980). Size composition, age composition, and growth of canary rockfish, *Sebastes pinniger*, and splitnose rockfish, *S. diploproa*, from the 1977 rockfish survey. *Marine Fisheries Review* **42**, 55–60.
- Boehlert, G. W., and Yoklavich, M. M. (1984). Variability in age estimates in *Sebastes* as a function of methodology, different readers, and different laboratories. *California Fish and Game* **70**, 210–224.
- Boehlert, G. W., and Yoklavich, M. M. (1986). Long-term cycles of growth in *Sebastes*: extracting information from otoliths. In 'Proceedings of the International Rockfish Symposium'. pp. 171–196. October 1986, Anchorage, Alaska, USA.
- Cailliet, G. M., Andrews, A. H., Burton, E. J., Watters, D. L., Kline, D. E., and Ferry-Graham, L. A. (2001). Age determination and validation studies of marine fishes: do deep-dwellers live longer? *Experimental Gerontology* **36**, 739–764. doi:10.1016/S0531-5565(00)00239-4
- Campana, S. E., Zwanenburg, K. C., and Smith, J. N. (1990). $^{210}\text{Pb}/^{226}\text{Ra}$ determination of longevity in redfish. *Canadian Journal of Fisheries and Aquatic Sciences* **47**, 163–165.
- Campana, S. E. (1997). Use of radiocarbon from nuclear fallout as a dated marker in the otoliths of haddock *Melanogrammus aeglefinus*. *Marine Ecology Progress Series* **150**, 49–56.
- Campana, S. E. (2001). Accuracy, precision and quality control in age determination, including a review of the use and abuse of age validation methods. *Journal of Fish Biology* **59**, 197–242. doi:10.1111/J.1095-8649.2001.TB00127.X
- Chilton, D. E., and Beamish, R. J. (1982). 'Age determination methods for fishes studied by the Groundfish Program at the Pacific Biological Station.' Canadian Special Publication of Fisheries and Aquatic Sciences No. 60. (Department of Fisheries and Oceans: Ottawa, Canada.)
- Crone, P. R., Piner, K. R., Methot, R. D., Conser, R. J., and Builder, T. L. (1999). 'Status of the canary rockfish resource off Oregon and Washington in 1999.' Appendix to 'The Status of the Pacific Coast Groundfish Fishery through 1999 and Recommended Acceptable Biological Catches for 2000'. (Pacific Fishery Management Council: Portland, OR.)
- DeMott, G. E. (1983). Movement of tagged lingcod and rockfishes off Depoe Bay, Oregon. M.S. Thesis. (Oregon State University, Portland, OR.)

- Francis, R. I. C. C. (2003). The precision of otolith radiometric ageing of fish and the effect of within-sample heterogeneity. *Canadian Journal of Fisheries and Aquatic Sciences* **60**, 441–447. doi:10.1139/F03-038
- Frantz, B. R., Foster, M. S., and Riosmena-Rodriguez, R. (2005). *Clathromorphum nereostratum* (Corallinales, Rhodophyta): the oldest alga? *Journal of Phycology* **41**, 770–773. doi:10.1111/J.1529-8817.2005.00107.X
- Gauldie, R. W., and Cremer, M. D. (2000). Confirmation of ^{222}Rn loss from otoliths of orange roughy *Hoplostethus atlanticus*. *Fisheries Science* **66**, 989–991. doi:10.1046/J.1444-2906.2000.00157.X
- Ingram, B. L., and Southon, J. R. (1996). Reservoir ages in eastern Pacific coastal and estuarine waters. *Radiocarbon* **38**, 572–582.
- Kalish, J. M. (1993). Pre- and post-bomb radiocarbon in fish otoliths. *Earth and Planetary Science Letters* **114**, 549–554. doi:10.1016/0012-821X(93)90082-K
- Kalish, J. M. (2001). 'Use of the Bomb Radiocarbon Chronometer to Validate Fish Age.' Final Report FRDC Project 93/109. (Fisheries Research and Development Corporation: Canberra.)
- Kalish, J. M., Nydal, R., Nedreaas, K. H., Burr, G. S., and Eine, G. L. (2001). A time history of pre- and post-bomb radiocarbon in the Barents Sea derived from Arcto-Norwegian cod otoliths. *Radiocarbon* **43**(2B), 843–855.
- Kastelle, C. R., Kimura, D. K., and Jay, S. R. (2000). Using $^{210}\text{Pb}/^{226}\text{Ra}$ disequilibrium to validate conventional ages in scorpaenids (genera *Sebastes* and *Sebastolobus*). *Fisheries Research* **46**, 299–312. doi:10.1016/S0165-7836(00)00155-7
- Kastelle, C. R., and Forsberg, J. E. (2002). Testing for loss of ^{222}Rn from Pacific halibut (*Hippoglossus stenolepis*) otoliths. *Fisheries Research* **57**, 93–98. doi:10.1016/S0165-7836(01)00339-3
- Kerr, L. A., Andrews, A. H., Frantz, B. R., Coale, K. H., Brown, T. A., and Cailliet, G. M. (2004). Radiocarbon in otoliths of yelloweye rockfish (*Sebastes ruberrimus*): a reference time series for the coastal waters of southeast Alaska. *Canadian Journal of Fisheries and Aquatic Sciences* **61**, 443–451. doi:10.1139/F04-009
- Kerr, L. A., Andrews, A. H., Frantz, B. R., Coale, K. H., Brown, T. A., Munk, K., and Cailliet, G. M. (2005). Age validation of quillback rockfish (*Sebastes maliger*) using bomb radiocarbon. *Fishery Bulletin* **103**(1), 97–107.
- Kimura, D. K., and Kastelle, C. R. (1995). Perspectives on the relationship between otolith growth and the conversion of isotope activity ratios to fish ages. *Canadian Journal of Fisheries and Aquatic Sciences* **52**, 2296–2303.
- Kline, D. E. (1996). Radiochemical age verification for two deep-sea rockfishes (*Sebastolobus altivelis* and *S. alascanus*). M.Sc. Thesis, California State University. (Moss Landing Marine Laboratories: San Jose, CA.)
- Love, M. S., Yoklavich, M., and Thorsteinson, L. (2002). 'The Rockfishes of the Northeast Pacific.' (University of California Press: Berkeley, CA.)
- MacLellan, S. E. (1997). 'How to Age Rockfish (*Sebastes*) using *S. alutus* as an Example – the Otolith Burnt Section Technique.' Canadian Technical Report of Fisheries and Aquatic Sciences 2146.
- Method, R., and Piner, K. (2002). Stock assessment of canary rockfish off California, Oregon and Washington in 2001. Appendix in 'Status of the Pacific Coast Groundfish Fishery through 2002; Stock Assessment and Fishery Evaluation. Vol. I'. (Pacific Fishery Management Council: Portland, OR.)
- Method, R. D., and Stewart, I. J. (2005). 'Status of the U.S. Canary Rockfish Resource in 2005.' (National Marine Fisheries Service, Northwest Fisheries Science Center: Seattle, WA.)
- Phillips, J. B. (1964). Life history studies on ten species of rockfish (genus *Sebastes*). *California Department of Fish and Game Fish Bulletin* **126**, 70 pp.
- Piner, K. R., and Wischniowski, S. G. (2004). Pacific halibut chronology of bomb radiocarbon in otoliths from 1941 to 1981 and a validation of ageing methods. *Journal of Fish Biology* **64**, 1060–1071. doi:10.1111/J.1095-8649.2004.0371.X
- Piner, K. R., Hamel, O. S., Henkel, J. L., Wallace, J. R., and Hutchinson, C. E. (2005). Age validation of canary rockfish (*Sebastes pinniger*) from off the Oregon coast (USA) using the bomb radiocarbon method. *Canadian Journal of Fisheries and Aquatic Sciences* **62**, 1060–1066. doi:10.1139/F05-082
- Six, L. D., and Horton, H. F. (1977). Analysis of age determination methods for yellowtail rockfish, canary rockfish, and black rockfish off Oregon. *Fishery Bulletin* **75**, 405–414.
- Smith, J. N., Nelson, R., and Campana, S. E. (1991). The use of Pb-210/Ra-226 and Th-228/Ra-228 disequilibria in the ageing of otoliths of marine fish. In 'Radionuclides in the Study of Marine Processes'. (Eds P. J. Kershaw and D. S. Woodhead.) pp. 350–359. (Elsevier Applied Science: New York.)
- Stanley, R. D. (1999). 'Canary Rockfish Assessment for 1999 and Recommended Yield Options for 2000/2001.' Canadian Stock Assessment Secretariat, Research Document 99/203. (Fisheries and Oceans Canada, Pacific Biological Station: Nanaimo, BC, Canada.)
- Stevens, M. M., Andrews, A. H., Cailliet, G. M., Coale, K. H., and Lundstrom, C. C. (2004). Radiometric validation of age, growth, and longevity for the blackgill rockfish, *Sebastes melanostomus*. *Fishery Bulletin* **102**, 711–722.
- Stuiver, M., and Polach, H. A. (1977). Discussion: reporting of ^{14}C data. *Radiocarbon* **19**, 355–363.
- Watters, D. L., Kline, D. E., Coale, K. H., and Cailliet, G. M. (2006). Radiometric age confirmation and growth of a deep-water marine fish species: the bank rockfish, *Sebastes rufus*. *Fisheries Research* **81**, 251–257. doi:10.1016/J.FISHRES.2006.06.019
- Westrheim, S. J., and Harling, W. R. (1975). 'Age-length Relationships for 26 Scorpaenids in the Northeastern Pacific Ocean.' Fisheries and Marine Service Research and Development Technical Report 565. (Research and Development Directorate, Pacific Biological Station: Nanaimo, BC, Canada.)
- Yamanaka, K. L., and Kronlund, A. R. (1997). 'Inshore Rockfish Stock Assessment for the West Coast of Canada in 1996 and Recommended Yields for 1997.' Canadian Technical Report of Fisheries and Aquatic Sciences No. 2175. (Department of Fisheries and Aquatic Sciences: Ottawa, Canada.)

Manuscript received 12 April 2007, accepted 11 May 2007